

Arterial segment	Median Bollinger score (IQR)		P-value*
	Non-diabetic	Diabetic	
Aorta	3 (1-3)	3 (3-3)	0.497
Common iliac	3 (2-7)	3 (1-3)	0.192
Internal iliac	3 (0-7)	3 (0-7)	0.994
External iliac	3 (0-7)	2 (0-3)	<0.05
Superficial femoral	8 (3-13)	7 (3-13)	0.411
<u>Profunda femoris</u>	0 (0-3)	0 (0-3)	0.889
Popliteal	5 (2-13)	6 (3-10)	0.369
Anterior tibial (proximal 3cm)	2 (0-8)	3 (0-7)	0.182
Anterior tibial (proximal 3 rd)†	3 (0-13)	5 (2-13)	0.109
Anterior tibial (middle 3 rd)†	3 (0-13)	4 (0-15)	0.112
Anterior tibial (distal 3 rd)†	3 (0-15)	7 (0-15)	0.417
Dorsalis pedis†	13 (0-15)	13 (0-15)	0.573
Peroneal (proximal 5cm)	3 (0-7)	3 (0-10)	0.386
Peroneal (proximal 3 rd)†	3 (0-7)	3 (0-13)	0.246
Peroneal (middle 3 rd)†	3 (0-7)	3 (0-13)	0.675
Peroneal (distal 3 rd)†	3 (0-13)	3 (0-15)	0.992
Posterior tibial (proximal 5cm)	3 (0-13)	6 (0-15)	<0.05
Posterior tibial (proximal 3 rd)†	3 (0-13)	8 (2-15)	<0.05
Posterior tibial (middle 3 rd)†	3 (0-15)	10 (0-15)	<0.05
Posterior tibial (distal 3 rd)†	3 (0-15)	13 (0-15)	<0.05
Medial plantar†	15 (0-15)	15 (0-15)	0.463
Lateral plantar†	15 (0-15)	15 (0-15)	0.978

* Mann-Whitney U Test

† Not originally included in Bollinger score

Table 1: Comparing the median Bollinger score of diabetics to non-diabetics by arterial segment